

Classification of indefinite hyper-Kähler symmetric spaces

Dmitri V. Alekseevsky¹, Vicente Cortés²

daleksee@mpim-bonn.mpg.de , vicente@math.uni-bonn.de ,

¹ Центр “Софус Ли”, Ген. Антонова 2 - 99, 117279 Москва and MPI für Mathematik, Bonn

² Mathematisches Institut der Universität Bonn, Beringstr. 1, D-53115 Bonn

Abstract

We classify indefinite simply connected hyper-Kähler symmetric spaces. Any such space without flat factor has commutative holonomy group and signature $(4m, 4m)$. We establish a natural 1-1 correspondence between simply connected hyper-Kähler symmetric spaces of dimension $8m$ and orbits of the group $GL(m, \mathbb{H})$ on the space $(S^4\mathbb{C}^n)^\tau$ of homogeneous quartic polynomials S in $n = 2m$ complex variables satisfying the reality condition $S = \tau S$, where τ is the real structure induced by the quaternionic structure of $\mathbb{C}^{2m} = \mathbb{H}^m$. We define and classify also complex hyper-Kähler symmetric spaces. Such spaces without flat factor exist in any (complex) dimension divisible by 4.

1 Introduction

We recall that a pseudo-Riemannian manifold (M, g) is called a symmetric space if any point $x \in M$ is an isolated fixed point of an involutive isometry s_x (called central symmetry with centre x). Since the product of two central symmetries s_x and s_y with sufficiently close centres is a shift along the geodesic (xy) , the group generated by central symmetries acts transitively on M and one can identify M with the quotient $M = G/K$ where G is the connected component of the isometry group $\text{Isom}(M, g)$ and K is the stabilizer of a point $o \in M$.

A symmetric space $(M = G/K, g)$ is called Kähler (respectively, hyper-Kähler) if its holonomy group $\text{Hol}(M, g)$ is a subgroup of the pseudo-unitary group $U(p, q)$ (respectively, of the pseudo-symplectic group $\text{Sp}(p, q) \subset \text{SU}(2p, 2q)$). Any hyper-Kähler symmetric space is in particular a homogeneous hypercomplex manifold. Homogeneous hypercomplex manifolds of compact Lie groups were constructed by Ph. Spindel, A. Sevrin, W. Troost, A. Van Proeyen [SSTVP] and by D. Joyce [J] and homogeneous hypercomplex structures on solvable Lie groups by M.L. Barberis and I. Dotti-Miatello [BD].

The classification of simply connected symmetric spaces reduces to the classification of involutive automorphisms σ of a Lie algebra $\mathfrak{g}$, such that the adjoint representation $\text{ad}_{\mathfrak{k}}|_{\mathfrak{m}}$ preserves a pseudo-Euclidean scalar product g , where

$$\mathfrak{g} = \mathfrak{k} + \mathfrak{m}, \quad \sigma|_{\mathfrak{k}} = 1, \quad \sigma|_{\mathfrak{m}} = -1,$$

is the eigenspace decomposition of the involution σ . Note that the eigenspace decomposition of an involutive automorphism is characterized by the conditions

$$[\mathfrak{k}, \mathfrak{k}] \subset \mathfrak{k}, \quad [\mathfrak{k}, \mathfrak{m}] \subset \mathfrak{m}, \quad [\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{k}.$$

Such a decomposition is called a symmetric decomposition.

In fact, for any pseudo-Riemannian symmetric space $M = G/K$ the conjugation with respect to the central symmetry s_o with centre $o = eK$ is an involutive automorphism of the Lie group G , which induces an involutive automorphism σ of its Lie algebra $\mathfrak{g}$. The pseudo-Riemannian metric of M induces a $\mathfrak{k}$ -invariant scalar product on $\mathfrak{m} \cong T_o M$, where $\mathfrak{g} = \mathfrak{k} + \mathfrak{m}$ is the symmetric decomposition defined by σ . Conversely, a symmetric decomposition $\mathfrak{g} = \mathfrak{k} + \mathfrak{m}$ together with a $\mathfrak{k}$ -invariant scalar product on $\mathfrak{m}$ determines a pseudo-Riemannian symmetric space $M = G/K$ where G is the simply connected Lie group with the Lie algebra $\mathfrak{g}$, K is the connected (and closed) subgroup of G generated by $\mathfrak{k}$, the pseudo-Riemannian metric on M is defined by g and the central symmetry is defined by the involutive automorphism σ associated to the symmetric decomposition.

Naturally identifying the space $\mathfrak{m}$ with the tangent space $T_o M$, the isotropy group is identified with $\text{Ad}_K|_{\mathfrak{m}}$ and the holonomy algebra is identified with $\text{ad}_{\mathfrak{h}}$ where $\mathfrak{h} = [\mathfrak{m}, \mathfrak{m}]$. If one assumes that the holonomy algebra is irreducible then one can prove that the Lie algebra $\mathfrak{g}$ is semisimple. Hence the classification of pseudo-Riemannian symmetric spaces with irreducible holonomy reduces to the classification of involutive automorphisms of semisimple Lie algebras. Such a classification was obtained by M. Berger [B1, B2] and A. Fedenko [F]. It includes the classification of Riemannian symmetric spaces (obtained earlier by E. Cartan), since according to de Rham's theorem any simply connected complete Riemannian manifold is a direct product of Riemannian manifolds with irreducible holonomy algebra and a Euclidean space.

A classification of pseudo-Riemannian symmetric spaces with non completely reducible holonomy is known only for signature $(1, n)$ (Cahen-Wallach [C-W]) and for signature $(2, n)$ under the assumption that the holonomy group is solvable (Cahen-Parker [C-P]). The classification problem for arbitrary signature looks very complicated and includes, for example, the classification of Lie algebras which admit a nondegenerate ad-invariant symmetric bilinear form. Some inductive construction of solvable Lie algebras with such a form was given by V. Kac.

In this paper we give a classification of pseudo-Riemannian hyper-Kähler symmetric spaces.

In particular, we prove that any simply connected hyper-Kähler symmetric space M has signature $(4m, 4m)$ and its holonomy group is commutative. The main result is the following, see Theorem 6.

Let (E, ω, j) be a complex symplectic vector space of dimension $4m$ with a quaternionic structure j such that $\omega(jx, jy) = \overline{\omega(x, y)}$ for all $x, y \in E$ and $E = E_+ \oplus E_-$ a j -invariant Lagrangian decomposition. Such a decomposition exists if and only if the Hermitian form $\gamma = \omega(\cdot, j\cdot)$ has real signature $(4m, 4m)$. We denote by τ the real structure in $S^{2r}E$ defined by $\tau(e_1 e_2 \dots e_{2r}) := j(e_1)j(e_2) \dots j(e_{2r})$, $e_i \in E$. Then any element $S \in (S^4 E_+)^{\tau}$ defines a hyper-Kähler symmetric space M_S which is associated with the symmetric decomposition

$$\mathfrak{g} = \mathfrak{h} + \mathfrak{m},$$

where $\mathfrak{m} = (\mathbb{C}^2 \otimes E)^{\rho}$ is the fixed point set of the real structure ρ on $\mathfrak{m}$ given by $\rho(h \otimes e) = j_H h \otimes j e$, where j_H is the standard quaternionic structure on $\mathbb{C}^2 = \mathbb{H}$, $\mathfrak{h} = \text{span}\{S_{ee'} | e, e' \in E\}^{\tau} \subset \text{sp}(E)^{\tau} \cong \text{sp}(m, m)$ with the natural action on $\mathfrak{m} \subset \mathbb{C}^2 \otimes E$ and the Lie bracket $\mathfrak{m} \wedge \mathfrak{m} \rightarrow \mathfrak{h}$ is given by

$$[h \otimes e, h' \otimes e'] = \omega_H(h, h') S_{e, e'}.$$

Moreover, we establish a natural 1-1 correspondence between simply connected hyper-Kähler symmetric spaces (up to isomorphism) and orbits of the group $GL(m, \mathbb{H})$ in $(S^4 E_+)^{\tau}$.

We define also the notion of complex hyper-Kähler symmetric space as a complex manifold (M, g) of complex dimension $4n$ with holomorphic metric g such that for any point $x \in M$ there is a holomorphic central symmetry s_x with centre x and which has holonomy group $\text{Hol}(M, g) \subset \text{Sp}(n, \mathbb{C})$ ($\text{Sp}(n, \mathbb{C}) \hookrightarrow \text{Sp}(n, \mathbb{C}) \times \text{Sp}(n, \mathbb{C}) \subset \text{O}(4n, \mathbb{C})$ is diagonally embedded) and give a classification of such spaces. We establish a natural 1-1 correspondence between simply connected complex hyper-Kähler symmetric spaces and homogeneous polynomials of degree 4 in the vector space $\mathbb{C}^{2n}$ considered up to linear transformations from $GL(2n, \mathbb{C})$.

2 Symmetric spaces

2.1 Basic facts about pseudo-Riemannian symmetric spaces

A **pseudo-Riemannian symmetric space** is a pseudo-Riemannian manifold (M, g) such that any point is an isolated fixed point of an isometric involution. Such a pseudo-Riemannian manifold admits a transitive Lie group of isometries L and can be identified with L/L_o , where L_o is the stabilizer of a point o . More precisely, any simply connected pseudo-Riemannian symmetric space $M = G/K$ is associated with a symmetric decomposition

$$\mathfrak{g} = \mathfrak{k} + \mathfrak{m}, \quad [\mathfrak{k}, \mathfrak{k}] \subset \mathfrak{k}, \quad [\mathfrak{k}, \mathfrak{m}] \subset \mathfrak{m}, \quad [\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{k} \quad (2.1)$$

of the Lie algebra $\mathfrak{g} = \text{Lie } G$ together with an Ad_K -invariant pseudo-Euclidean scalar product on $\mathfrak{m}$. We will assume that G acts almost effectively on M , i.e. $\mathfrak{k}$ does not contain any nontrivial ideal of $\mathfrak{g}$, that M and G are simply connected and that K is connected. Then, under the natural identification of the tangent space $T_o M$ at the canonical base point $o = eK$ with $\mathfrak{m}$, the holonomy group $\text{Hol} \subset \text{Ad}_K|_{\mathfrak{m}}$. We will denote by $\mathfrak{h}$ the holonomy Lie algebra. Since the isotropy representation is faithful it is identified with the subalgebra $\mathfrak{h} = [\mathfrak{m}, \mathfrak{m}] := \text{span}\{[x, y] | x, y \in \mathfrak{m}\} \subset \mathfrak{k}$. Recall that the curvature tensor R of a symmetric space M at o is $\mathfrak{h}$ -invariant and determines the Lie bracket in the ideal $\mathfrak{h} + \mathfrak{m} \subset \mathfrak{g}$ as follows:

$$\mathfrak{h} = R(\mathfrak{m}, \mathfrak{m}) := \text{span}\{R(x, y) | x, y \in \mathfrak{m}\} \quad \text{and} \quad [x, y] = -R(x, y), \quad x, y \in \mathfrak{m}.$$

The following result is well known:

Proposition 1 *The full Lie algebra of Killing fields of a symmetric space has the form*

$$\text{isom}(M) = \tilde{\mathfrak{h}} + \mathfrak{m},$$

where the full isotropy subalgebra is given by

$$\tilde{\mathfrak{h}} = \text{aut}(R) = \{A \in \text{so}(\mathfrak{m}) | A \cdot R = [A, R(\cdot, \cdot)] - R(A\cdot, \cdot) - R(\cdot, A\cdot) = 0\}. \quad (2.2)$$

2.2 Symmetric spaces of semisimple Lie groups

We will prove that in the case when $(M = G/K, g)$ is a pseudo-Riemannian symmetric space of a (connected) semisimple Lie group G then G is the maximal connected Lie group of isometries of M .

Proposition 2 *Let $(M = G/K, g)$ be a pseudo-Riemannian symmetric space associated with a symmetric decomposition $\mathfrak{g} = \mathfrak{k} + \mathfrak{m}$. If G is semisimple and almost effective then*

- (i) *the restriction of the Cartan-Killing form B of $\mathfrak{g}$ to $\mathfrak{k}$ is nondegenerate and hence $\mathfrak{k}$ is a reductive subalgebra of $\mathfrak{g}$ and $\mathfrak{g} = \mathfrak{k} + \mathfrak{m}$ is a B -orthogonal decomposition,*
- (ii) *$\mathfrak{k} = [\mathfrak{m}, \mathfrak{m}]$ and*
- (iii) *$\mathfrak{g} = \text{isom}(M, g)$ is the Lie algebra of the full isometry group of M .*

Proof: For (i) see [O-V] Ch. 3 Proposition 3.6.

(ii) It is clear that $\bar{\mathfrak{g}} = [\mathfrak{m}, \mathfrak{m}] + \mathfrak{m}$ is an ideal of $\mathfrak{g}$. The B -orthogonal complement $\mathfrak{a} := \bar{\mathfrak{g}}^\perp \subset \mathfrak{k}$ is a complementary ideal of $\mathfrak{g}$. Since $[\mathfrak{a}, \mathfrak{m}] = 0$ the Lie algebra $\mathfrak{a}$ acts trivially on M . From the effectivity of $\mathfrak{g}$ we conclude that $\mathfrak{a} = 0$.

(iii) By Proposition 1, $\tilde{\mathfrak{g}} = \text{isom}(M, g) = \tilde{\mathfrak{h}} + \mathfrak{m}$, where $\tilde{\mathfrak{h}} = \text{aut}(R) = \{A \in \text{so}(\mathfrak{m}) | A \cdot R = 0\}$. Now $\tilde{\mathfrak{h}}$ preserves $\mathfrak{m}$ and by the identity $A \cdot R = [A, R(\cdot, \cdot)] - R(A \cdot, \cdot) - R(\cdot, A \cdot)$ it also normalizes $\mathfrak{k}$. This shows that $\tilde{\mathfrak{h}}$ normalizes $\mathfrak{g}$ and hence $\mathfrak{g} \subset \tilde{\mathfrak{g}}$ is an ideal. Since $\mathfrak{g}$ is semisimple there exists a $\mathfrak{g}$ -invariant complement $\mathfrak{b}$ in $\tilde{\mathfrak{g}}$. Note that $[\mathfrak{g}, \mathfrak{b}] \subset \mathfrak{g} \cap \mathfrak{b} = 0$. We can decompose any $X \in \mathfrak{b}$ as $X = Y + Z$, where $Y \in \tilde{\mathfrak{h}}$ and $Z \in \mathfrak{m}$. From $[\mathfrak{g}, \mathfrak{b}] = 0$ it follows that $[\mathfrak{g}, Y] = [\mathfrak{g}, Z] = 0$ and in particular $[\mathfrak{m}, Y] = 0$. This implies that $Y = 0$ and $X = Z \in \mathfrak{b} \cap \mathfrak{m} = 0$. This shows that $\mathfrak{b} = 0$ proving (iii). $\square$

We recall that a pseudo-Riemannian Hermitian symmetric space is pseudo-Riemannian symmetric space $(M = G/K, g)$ together with an invariant (and hence parallel) g -orthogonal complex structure J .

Proposition 3 *Let $(M = G/K, g, J)$ be a pseudo-Riemannian Hermitian symmetric space of a semisimple and almost effective Lie group G . Then the Ricci curvature of M is not zero.*

Proof: From Proposition 2 it follows that $\mathfrak{g} = \text{isom}(M, g) = \tilde{\mathfrak{h}} + \mathfrak{m}$, where $\tilde{\mathfrak{h}} = \mathfrak{k} = [\mathfrak{m}, \mathfrak{m}]$. It is well known that the curvature tensor R of any pseudo-Kähler manifold (and in particular of any pseudo-Riemannian Hermitian symmetric space) is invariant under the operator J . This shows that $J \in \tilde{\mathfrak{h}} = \text{aut}(R) = [\mathfrak{m}, \mathfrak{m}] = \mathfrak{h}$ (holonomy Lie algebra), which implies that the holonomy Lie algebra is not a subalgebra of $\text{su}(\mathfrak{m}) \cong \text{su}(p, q)$. Hence M is not Ricci-flat. In fact, we can write $J = \sum \text{ad}[X_i, Y_i]$, for $X_i, Y_i \in \mathfrak{m}$. Then using the formulas $\text{Ric}(X, JY) = \text{tr} J R(X, Y)$ for the Ricci curvature of a pseudo-Kähler manifold and $R(X, Y) = -\text{ad}_{[X, Y]}|_{\mathfrak{m}}$ for the curvature of a symmetric space we calculate:

$$\sum \text{Ric}(X_i, JY_i) = \sum \text{tr} J R(X_i, Y_i) = - \sum \text{tr} J \text{ad}[X_i, Y_i] = -\text{tr} J^2 \neq 0.$$

$\square$

3 Structure of hyper-Kähler symmetric spaces

3.1 Definitions

A (possibly indefinite) **hyper-Kähler manifold** is a pseudo-Riemannian manifold (M^{4n}, g) of signature $(4k, 4l)$ together with a **compatible** hypercomplex structure, i.e. three g -orthogonal parallel complex structures $(J_1, J_2, J_3 = J_1 J_2)$. This means that the holonomy group $\text{Hol} \subset \text{Sp}(k, l)$. Two hyper-Kähler manifolds (M, g, J_α) ($\alpha = 1, 2, 3$) and (M', g', J'_α) are called **isomorphic** if there exists a triholomorphic isometry $\varphi : M \rightarrow M'$, i.e. $\varphi^* J'_\alpha = J_\alpha$ and $\varphi^* g' = g$.

A **hyper-Kähler symmetric space** is a pseudo-Riemannian symmetric space $(M = G/K, g)$ together with an invariant compatible hypercomplex structure. Consider now a simply con-

nected hyper-Kähler symmetric space $(M = G/K, g, J_\alpha)$. Without restriction of generality we will assume that G acts almost effectively. M being hyper-Kähler is equivalent to $\text{Ad}_K|_{\mathfrak{m}} \subset \text{Sp}(k, l)$, or, since K is connected, to $\text{ad}_{\mathfrak{k}}|_{\mathfrak{m}} \subset \text{sp}(k, l)$. This condition means that $\mathfrak{k}$ commutes with the Lie algebra $Q = \text{sp}(1) \subset \text{so}(\mathfrak{m}) = \text{so}(4k, 4l)$ spanned by three anticommuting complex structures J_1, J_2, J_3 .

3.2 Existence of a transitive solvable group of isometries and solvability of the holonomy

In this subsection we prove that any simply connected hyper-Kähler symmetric space (M, g, J_α) admits a transitive solvable Lie group $G \subset \text{Aut}(g, J_\alpha)$ of automorphisms and has solvable holonomy group.

Proposition 4 *Let $(M = G/K, g, J_\alpha)$ be a simply connected hyper-Kähler symmetric space and $A = \text{Aut}_0(g, Q) \supset \text{Aut}_0(g, J_\alpha) \supset G$ the connected group of isometries which preserve the quaternionic structure $Q = \text{span}\{J_\alpha\}$. Then*

- (i) *the stabilizer A_o of a point $o \in M$ contains a maximal semisimple subgroup of A ,*
- (ii) *the radical R of A acts transitively and triholomorphically on M and*
- (iii) *the holonomy group of M is solvable.*

Proof: We consider the quaternionic Kähler symmetric space $(M = A/A_o, g, Q)$. The Lie algebra $\mathfrak{a}_o$ of the stabilizer is given by

$$\mathfrak{a}_o = \text{aut}(R, Q) = \{A \in \text{so}(T_o M) | A \cdot R = 0, [A, Q] \subset Q\}.$$

Since the curvature tensor of a quaternionic Kähler manifold is invariant under the quaternionic structure Q we conclude that $Q \subset \mathfrak{a}_o$ and $\mathfrak{a}_o = Q \oplus Z_{\mathfrak{a}}(Q)$, where $Z_{\mathfrak{a}}(Q)$ denotes the centralizer of Q in $\mathfrak{a}$. Since $Q \cong \text{sp}(1)$ is simple, we may choose a Levi-Malcev decomposition $\mathfrak{a} = \mathfrak{s} + \mathfrak{r}$ such that the Levi subalgebra $\mathfrak{s} \supset Q$. We put $\mathfrak{m}_r := [Q, \mathfrak{r}]$ and denote by $\mathfrak{m}_s$ a $Q \oplus Z_{\mathfrak{s}}(Q)$ -invariant complement of Q in $[Q, \mathfrak{s}]$. The stabilizer has the decomposition $\mathfrak{a}_o = Q \oplus (Z_{\mathfrak{s}}(Q) + Z_{\mathfrak{r}}(Q))$.

Lemma 1 *The complement $\mathfrak{m} = \mathfrak{m}_s + \mathfrak{m}_r$ to $\mathfrak{a}_o$ in $\mathfrak{a}$ is $\mathfrak{a}_o$ -invariant and the decomposition*

$$\mathfrak{a} = \mathfrak{a}_o + \mathfrak{m}$$

is a symmetric decomposition.

Proof: It is clear that $\mathfrak{m}_r$ is $\mathfrak{a}_o$ -invariant and $\mathfrak{m}_s$ is invariant under $Q \oplus Z_{\mathfrak{s}}(Q)$ by construction. It remains to check that $[Z_{\mathfrak{r}}(Q), \mathfrak{m}_s] \subset \mathfrak{m}$. Since $\mathfrak{m}_s = [Q, \mathfrak{m}_s]$, we have

$$[Z_{\mathfrak{r}}(Q), \mathfrak{m}_s] = [Z_{\mathfrak{r}}(Q), [Q, \mathfrak{m}_s]] = [Q, [Z_{\mathfrak{r}}(Q), \mathfrak{m}_s]] \subset [Q, \mathfrak{r}] = \mathfrak{m}_r \subset \mathfrak{m}.$$

This shows that $\mathfrak{a} = \mathfrak{a}_o + \mathfrak{m}$ is an $\mathfrak{a}_o$ -invariant decomposition. We denote by $\mathfrak{a} = \mathfrak{a}_o + \mathfrak{p}$ a symmetric decomposition. Any other $\mathfrak{a}_o$ -invariant decomposition is of the form $\mathfrak{a} = \mathfrak{a}_o + \mathfrak{p}_\varphi$, where $\varphi : \mathfrak{p} \rightarrow \mathfrak{a}_o$ is an $\mathfrak{a}_o$ -equivariant map and $\mathfrak{p}_\varphi = \{X + \varphi(X) | X \in \mathfrak{p}\}$. If such non-zero equivariant map φ exists then $\mathfrak{p}$ and $\mathfrak{a}_o$ contain non-trivial isomorphic Q -submodules. Since $\mathfrak{p}$ is a sum of 4-dimensional irreducible Q -modules and $\mathfrak{a}_o$ is the sum of the 3-dimensional irreducible Q -module Q and the trivial complementary Q -module $Z_{\mathfrak{a}_o}(Q)$, we infer that there exists a unique $\mathfrak{a}_o$ -invariant decomposition, which coincides with the symmetric decomposition $\mathfrak{a} = \mathfrak{a}_o + \mathfrak{p}$. $\square$

To prove (i) we have to check that $\mathfrak{m}_s = 0$. We note that by the previous lemma $\mathfrak{s} = (Q \oplus Z_{\mathfrak{s}}(Q)) + \mathfrak{m}_s$ is a symmetric decomposition of the semisimple Lie algebra $\mathfrak{s}$. Since $[\mathfrak{m}_s, \mathfrak{m}_s] \subset Z_{\mathfrak{s}}(Q)$ it defines a hyper-Kähler symmetric space $N = L/L_o$, where L is the simply connected semisimple Lie group with Lie algebra $\mathfrak{l} = Z_{\mathfrak{s}}(Q) + \mathfrak{m}_s$ and L_o is the Lie subgroup generated by the subalgebra $Z_{\mathfrak{s}}(Q) \subset \mathfrak{l}$. Since N is in particular a Ricci-flat pseudo-Riemannian Hermitian symmetric space, from Proposition 3 we obtain that N is reduced to point. Therefore $\mathfrak{m}_s = 0$. This proves (i) and (ii). Finally, since the holonomy Lie algebra $\mathfrak{h}$ is identified with $\mathfrak{h} = [\mathfrak{m}, \mathfrak{m}] = [\mathfrak{m}_r, \mathfrak{m}_r] \subset \mathfrak{r}$ it is solvable as subalgebra of the solvable Lie algebra $\mathfrak{r}$. $\square$

3.3 Hyper-Kähler symmetric spaces and second prolongation of symplectic Lie algebras

Let $(M = G/K, g, J_\alpha)$ be a simply connected hyper-Kähler symmetric space associated with a symmetric decomposition (2.1). Without restriction of generality we will assume that G acts almost effectively and that $\mathfrak{k} = [\mathfrak{m}, \mathfrak{m}] = \mathfrak{h}$ (holonomy Lie algebra). The complexification $\mathfrak{m}^{\mathbb{C}}$ as $\mathfrak{h}^{\mathbb{C}}$ -module can be written as $\mathfrak{m}^{\mathbb{C}} = H \otimes E$, such that $\mathfrak{h}^{\mathbb{C}} \subset \text{Id} \otimes \mathfrak{sp}(E) \cong \mathfrak{sp}(E)$, where $H = \mathbb{C}^2$ and $E = \mathbb{C}^{2n}$ are complex symplectic vector spaces with symplectic form ω_H and ω_E , respectively, such that $g^{\mathbb{C}} = \omega_H \otimes \omega_E$ is the complex bilinear metric on $\mathfrak{m}^{\mathbb{C}}$ induced by g . Note that the symplectic forms are unique up to the transformation $\omega_H \mapsto \lambda \omega_H$, $\omega_E \mapsto \lambda^{-1} \omega_E$, $\lambda \in \mathbb{C}^*$. We have also quaternionic structures j_H and j_E on H and E , such that $\omega_H(j_H x, j_H y) = \overline{\omega_H(x, y)}$ for all $x, y \in H$ and $\omega_E(j_E x, j_E y) = \overline{\omega_E(x, y)}$ for all $x, y \in E$, where the bar denotes complex conjugation. This implies that $\gamma_H := \omega_H(\cdot, j_H \cdot)$ and $\gamma_E := \omega_E(\cdot, j_E \cdot)$ are Hermitian forms on H and E . For fixed ω_H and ω_E the quaternionic structures j_H and j_E are uniquely determined if we require that γ_H is positive definite and that $\rho = j_H \otimes j_E$ is the real structure on $\mathfrak{m}^{\mathbb{C}}$, i.e. the complex conjugation with respect to $\mathfrak{m}$. The metric $g^{\mathbb{C}}$

and the Hermitian form $g^{\mathbb{C}}(\cdot, \rho \cdot) = \gamma_H \otimes \gamma_E$ restrict to a real valued scalar product g of some signature $(4k, 4l)$ on $\mathfrak{m} = (H \otimes E)^\rho$, where $(2k, 2l)$ is the (real) signature of the Hermitian form $\gamma_E = \omega_E(\cdot, j_E \cdot)$. Note that for the holonomy algebra we have the inclusion

$$\mathfrak{h} = \text{Id} \otimes (\mathfrak{h}^{\mathbb{C}})^{j_E} \hookrightarrow \text{sp}(E)^{j_E} = \{A \in \text{sp}(E) \mid [A, j_E] = 0\} = \text{aut}(E, \omega_E, j_E) \cong \text{aut}(\mathfrak{m}, g, J_\alpha) \cong \text{sp}(k, l).$$

Using the symplectic forms we identify $H = H^*$ and $E = E^*$. Then the symplectic Lie algebras are identified with symmetric tensors as follows:

$$\text{sp}(H) = S^2 H, \quad \text{sp}(E) = S^2 E.$$

Since the curvature tensor R of any hyper-Kähler manifold M^{4n} can be identified at a point $p \in M$ with an element $R \in S^2 \text{sp}(k, l)$ it is invariant under the Lie algebra $\text{sp}(1) = \text{span}\{J_1, J_2, J_3\}$. Let $M = G/K$ be a hyper-Kähler symmetric space as above. By Proposition 1 we can extend the Lie algebra $\mathfrak{g} = \mathfrak{k} + \mathfrak{m} = \mathfrak{h} + \mathfrak{m}$ to a Lie algebra

$$\tilde{\mathfrak{g}} = \text{sp}(1) + \mathfrak{h} + \mathfrak{m}$$

of Killing vector fields such that $[\text{sp}(1), \mathfrak{h}] = 0$. In the $H \otimes E$ -formalism the Lie algebra $\text{sp}(1)$ is identified with $\text{sp}(H)^{j_H} \otimes \text{Id} \subset \text{so}(\mathfrak{m})$.

Lemma 2 *Denote by $\tilde{\mathfrak{g}}^{\mathbb{C}} = \text{sp}(1, \mathbb{C}) + \mathfrak{h}^{\mathbb{C}} + \mathfrak{m}^{\mathbb{C}}$ the complexification of the Lie algebra $\tilde{\mathfrak{g}}$. Then the Lie bracket $[\cdot, \cdot] : \wedge^2 \mathfrak{m}^{\mathbb{C}} \rightarrow \mathfrak{h}^{\mathbb{C}}$ can be written as*

$$[h \otimes e, h' \otimes e'] = \omega_H(h, h') S_{e, e'}, \quad (3.1)$$

where $S \in (\mathfrak{h}^{\mathbb{C}})^{(2)} := \mathfrak{h}^{\mathbb{C}} \otimes S^2 E^* \cap E \otimes S^3 E^* = \mathfrak{h}^{\mathbb{C}} \otimes \mathfrak{h}^{\mathbb{C}} \cap S^4 E$. Moreover S is $\text{sp}(1, \mathbb{C}) \oplus \mathfrak{h}^{\mathbb{C}}$ -invariant and satisfies the following reality condition: $[S_{j_E e, e'} - S_{e, j_E e'}, j_E] = 0$.

Proof: The Lie bracket $[\cdot, \cdot] : \wedge^2 \mathfrak{m}^{\mathbb{C}} \rightarrow \mathfrak{h}^{\mathbb{C}}$ is an $\text{sp}(1, \mathbb{C}) \oplus \mathfrak{h}^{\mathbb{C}}$ -equivariant map, due to the Jacobi identity. We decompose the $\text{sp}(H) \oplus \text{sp}(E)$ -module $\wedge^2 \mathfrak{m}^{\mathbb{C}}$:

$$\wedge^2 \mathfrak{m}^{\mathbb{C}} = \wedge^2 (H \otimes E) = \wedge^2 H \otimes S^2 E \oplus S^2 H \otimes \wedge^2 E = \omega_H \otimes S^2 E \oplus S^2 H \otimes \wedge^2 E.$$

Since $\mathfrak{h}^{\mathbb{C}} \subset S^2 E$ the Lie bracket defines an $\text{sp}(1, \mathbb{C}) \oplus \mathfrak{h}^{\mathbb{C}}$ -invariant element of the space $\omega_H \otimes S^2 E \otimes S^2 E \oplus S^2 H \otimes \wedge^2 E \otimes S^2 E$. The second summand has no nontrivial $\text{sp}(1, \mathbb{C})$ -invariant elements. Hence the bracket is of the form (3.1), where $S \in S^2 E^* \otimes \mathfrak{h}^{\mathbb{C}} \subset S^2 E \otimes S^2 E$. The Jacobi identity reads:

$$\begin{aligned} 0 &= [h \otimes e, [h' \otimes e', h'' \otimes e'']] - [[h \otimes e, h' \otimes e'], h'' \otimes e''] - [h' \otimes e', [h \otimes e, h'' \otimes e'']] \\ &= -\omega_H(h', h'') h \otimes S_{e', e''} e - \omega_H(h, h') h'' \otimes S_{e, e'} e'' + \omega_H(h, h'') h' \otimes S_{e, e''} e'. \end{aligned}$$

Since $\dim H = 2$ we may assume that $h, h' = h''$ is a symplectic basis, i.e. $\omega_H(h, h') = 1$, and the equation implies: $S_{e, e''} e' = S_{e, e'} e''$, i.e. $S \in (\mathfrak{h}^\mathbb{C})^{(2)}$. The Lie bracket of two real elements $h \otimes e + j_H h \otimes j_E e$ and $h \otimes e' + j_H h \otimes j_E e' \in \mathfrak{m} \subset \mathfrak{m}^\mathbb{C}$ is an element of $\mathfrak{h}$. This gives:

$$[h \otimes e + j_H h \otimes j_E e, h \otimes e' + j_H h \otimes j_E e'] = \omega_H(h, j_H h)(S_{e, j_E e'} - S_{j_E e, e'}) \in \mathfrak{h}.$$

From the fact that the Hermitian form $\gamma_H = \omega_H(\cdot, j_H \cdot)$ is positive definite it follows that $\omega_H(h, j_H h) \neq 0$. This establishes the reality condition since $\mathfrak{h} = \{A \in \mathfrak{h}^\mathbb{C} | [A, j_E] = 0\}$. $\square$

In fact any tensor $S \in S^4 E$ satisfying the conditions of the above lemma can be used to define a hyper-Kähler symmetric space as the following theorem shows. We can identify $S^4 E$ with the space $\mathbb{C}[E]^{(4)}$ of homogeneous quartic polynomials on $E \cong E^*$.

Theorem 1 *Let $S \in S^4 E$, $E = \mathbb{C}^{2n}$, be a quartic polynomial invariant under all endomorphisms $S_{e, e'} \in S^2 E = \text{sp}(E)$ and satisfying the reality condition*

$$[S_{j_E e, e'} - S_{e, j_E e'}, j_E] = 0. \quad (3.2)$$

Then it defines a hyper-Kähler symmetric space, which is associated with the following complex symmetric decomposition

$$\mathfrak{g}^\mathbb{C} = \mathfrak{h}^\mathbb{C} + H \otimes E, \quad \mathfrak{h}^\mathbb{C} = \text{span}\{S_{e, e'} | e, e' \in E\} \subset \text{sp}(E). \quad (3.3)$$

The bracket $\wedge^2(H \otimes E) \rightarrow \mathfrak{h}^\mathbb{C}$ is given by (3.1). The real symmetric decomposition is defined as ρ -real form $\mathfrak{g} = \mathfrak{h} + \mathfrak{m}$ of (3.3), where

$$\mathfrak{k} = \mathfrak{h} = \{A \in \mathfrak{h}^\mathbb{C} | [A, j_E] = 0\} = \text{span}\{S_{j_E e, e'} - S_{e, j_E e'} | e, e' \in E\}, \quad \mathfrak{m} = (H \otimes E)^\rho.$$

The hyper-Kähler symmetric space M associated to this symmetric decomposition is the quotient $M = M_S = G/K$, where G is the simply connected Lie group with Lie algebra $\mathfrak{g}$ and $K \subset G$ is the connected (and closed) subgroup with Lie algebra $\mathfrak{k} = \mathfrak{h}$.

Moreover any simply connected hyper-Kähler symmetric space can be obtained by this construction. Two hyper-Kähler symmetric spaces M_S and $M_{S'}$ defined by quartics S and S' are isomorphic if and only if S and S' are in the same orbit of the group $\text{Aut}(E, \omega_E, j_E) = \{A \in \text{Sp}(E) | [A, j_E] = 0\} \cong \text{Sp}(k, l)$.

Proof: First of all we note that $\mathfrak{h}^\mathbb{C} = S_{E, E} := \text{span}\{S_{e, e'} | e, e' \in E\}$ is a subalgebra of $\text{sp}(E)$ because

$$[S_{e, e'}, S_{f, f'}] = (S_{e, e'} \cdot S)_{f, f'} - S_{S_{e, e'} f, f'} - S_{f, S_{e, e'} f'} = -S_{S_{e, e'} f, f'} - S_{f, S_{e, e'} f'} \in \mathfrak{h}^\mathbb{C}.$$

Since S is $\mathfrak{h}^\mathbb{C}$ -invariant and completely symmetric we can check, as in Lemma 2, that the Jacobi identity is satisfied and that (3.3) defines a complex symmetric decomposition. We prove that $\mathfrak{h} := \text{span}\{S_{j_E e, e'} - S_{e, j_E e'} | e, e' \in E\} \subset \{A \in \mathfrak{h}^\mathbb{C} | [A, j_E] = 0\}$ defines a real form of $\mathfrak{h}^\mathbb{C}$. Indeed for $e, e' \in E$ we have

$$\begin{aligned} S_{e, e'} &= \frac{1}{2}(S_{e, e'} + S_{j_E e, j_E e'}) - \frac{\sqrt{-1}}{2}(\sqrt{-1}S_{e, e'} - \sqrt{-1}S_{j_E e, j_E e'}) \\ &= \frac{1}{2}(S_{j_E e'', e'} - S_{e'', j_E e'}) - \frac{\sqrt{-1}}{2}(S_{j_E e'', \sqrt{-1}e'} - S_{e'', j_E \sqrt{-1}e'}), \end{aligned}$$

where $e'' = -j_E e$. Due to the reality condition the restriction of the Lie bracket $[\cdot, \cdot] : \wedge^2 \mathfrak{m}^\mathbb{C} \rightarrow \mathfrak{h}^\mathbb{C}$ to $\wedge^2 \mathfrak{m}$ has values in $\mathfrak{h}$ and $\mathfrak{g} = \mathfrak{h} + \mathfrak{m}$ is a symmetric decomposition with $[\mathfrak{m}, \mathfrak{m}] = \mathfrak{h}$. The metric $g^\mathbb{C} = \omega_H \otimes \omega_E$ defines a real valued scalar product g of some signature (p, q) on $\mathfrak{m} = (H \otimes E)^\rho$, which is invariant under the Lie algebra $\mathfrak{h}$. Since $[\mathfrak{h}, j_E] = 0$ the holonomy algebra $\mathfrak{h} \subset \text{sp}(k, l)$, $p = 4k$, $q = 4l$. Hence this symmetric decomposition defines a hyper-Kähler symmetric space.

By Lemma 2 any hyper-Kähler symmetric space can be obtained by this construction. It is well known that a simply connected symmetric space M of signature (p, q) is determined by its abstract curvature tensor $R \in S^2(\wedge^2 V)$, $V = \mathbb{R}^{p, q}$, and two tensors R and R' define isometric symmetric spaces if and only if they belong to the same $O(V)$ orbit. Similarly a simply connected hyper-Kähler symmetric space is determined up to isometry by its abstract curvature tensor $R \in S^2(\wedge^2 V)$, where $V = \mathbb{R}^{4k, 4l}$ is the pseudo-Euclidean vector space with fixed hypercomplex structure $J_\alpha \in O(V)$. For a hyper-Kähler symmetric space the complexified curvature tensor has the form

$$R(h \otimes e, h' \otimes e') = -\omega_H(h, h')S_{e, e'}.$$

where $S \in S^4 E$ is the quartic of Lemma 2. Two such curvature tensors define isomorphic hyper-Kähler symmetric spaces if and only if they belong to the same orbit of $\text{Aut}(\mathbb{R}^{4k, 4l}, J_\alpha) = \text{Sp}(k, l)$. The group $\text{Sp}(k, l)$ acts on $V^\mathbb{C} = H \otimes E$ as $\text{Id} \otimes \text{Sp}(E)^{j_E} = \text{Id} \otimes \text{Aut}(E, \omega_E, j_E)$. Hence two curvature tensors $R = -\omega_H \otimes S$ and $R' = -\omega_H \otimes S'$ are in the same $\text{Sp}(k, l)$ -orbit if and only if S and S' are in the same $\text{Sp}(k, l)$ -orbit on $S^4 E$. $\square$

4 Complex hyper-Kähler symmetric spaces

4.1 Complex hyper-Kähler manifolds

A **complex Riemannian manifold** is a complex manifold M equipped with a complex metric g , i.e. a holomorphic section $g \in \Gamma(S^2 T^* M)$ which defines a nondegenerate complex quadratic

form. As in the real case any such manifold has a unique holomorphic torsionfree and metric connection (Levi-Civita connection). A **complex hyper-Kähler manifold** is a complex Riemannian manifold (M^{4n}, g) of complex dimension $4n$ together with a compatible hypercomplex structure, i.e. three g -orthogonal parallel complex linear endomorphisms $(J_1, J_2, J_3 = J_1 J_2)$ with $J_\alpha^2 = -1$. This means that the holonomy group $\text{Hol} \subset \text{Sp}(n, \mathbb{C}) = Z_{O(4n, \mathbb{C})}(\text{Sp}(1, \mathbb{C}))$. The linear group $\text{Sp}(n, \mathbb{C})$ is diagonally embedded into $\text{Sp}(n, \mathbb{C}) \times \text{Sp}(n, \mathbb{C}) \subset \text{GL}(4n, \mathbb{C})$. Two complex hyper-Kähler manifolds (M, g, J_α) ($\alpha = 1, 2, 3$) and (M', g', J'_α) are called **isomorphic** if there exists a holomorphic isometry $\varphi : M \rightarrow M'$ such that $\varphi^* J'_\alpha = J_\alpha$ and $\varphi^* g' = g$.

We will show that the complex hyper-Kähler structure can be described as a half-flat Grassmann structure of a certain type. A **Grassmann structure** on a complex Riemannian manifold (M, g) is a decomposition of the (holomorphic) tangent bundle $TM \cong H \otimes E$ into the tensor product of two holomorphic vector bundles H and E of rank $2m$ and $2n$ with holomorphic nondegenerate 2-forms ω_H and ω_E such that $g = \omega_H \otimes \omega_E$. The Grassmann structure will be called **parallel** if the Levi-Civita connection $\nabla = \nabla^{TM}$ can be decomposed as:

$$\nabla = \nabla^H \otimes \text{Id} + \text{Id} \otimes \nabla^E,$$

where ∇^H and ∇^E are (uniquely defined) symplectic connections in the bundles H and E . A parallel Grassmann structure will be called **half-flat** if ∇^H is flat. Note that a parallel Grassmann structure on a simply connected manifold is half-flat if and only if the holonomy group of the Levi-Civita connection is contained in $\text{Id} \otimes \text{Sp}(n, \mathbb{C}) \subset \text{Sp}(m, \mathbb{C}) \otimes \text{Sp}(n, \mathbb{C}) \subset \text{O}(\mathbb{C}^{2m} \otimes \mathbb{C}^{2n})$.

Proposition 5 *A complex hyper-Kähler structure (g, J_α) on a simply connected complex manifold M is equivalent to the following geometric data:*

- (i) *a half-flat Grassmann structure $(TM, g, \nabla) \cong (H, \omega_H, \nabla^H) \otimes (E, \omega_E, \nabla^E)$ and*
- (ii) *an isomorphism of flat symplectic vector bundles $H \cong M \times \mathbb{C}^2$. Under this isomorphism $\omega_H = h_1^* \wedge h_2^*$, where (h_1, h_2) is the standard basis of $\mathbb{C}^2$ considered as parallel frame of the trivial bundle $H = M \times \mathbb{C}^2$.*

More precisely,

$$J_1 = R_i \otimes \text{Id}, \quad J_2 = R_j \otimes \text{Id}, \quad \text{and} \quad J_3 = R_k \otimes \text{Id}, \quad (4.1)$$

where we have identified $\mathbb{C}^2 = \mathbb{C}h_1 \oplus \mathbb{C}h_2$ with $\mathbb{H} = \text{span}_{\mathbb{R}}\{1, i, j, k\} = \text{span}_{\mathbb{C}}\{1, j\} = \mathbb{C}1 \oplus \mathbb{C}j$ with the complex structure defined by left-multiplication by i and R_x denotes the right-multiplication by the quaternion $x \in \mathbb{H}$.

Proof: It is easy to check that the geometric data (i) and (ii) define a complex hyper-Kähler structure on M . Conversely let (g, J_α) be a complex hyper-Kähler structure on M . The endomorphism J_1 has eigenvalues $\pm i$ and the tangent space can be decomposed into a sum of eigenspaces

$$TM = E_+ \oplus E_-.$$

From the J_1 -invariance of the metric g it follows that $g(E_\pm, E_\pm) = 0$ and we can identify $E_- = E^*$ with the dual space of $E = E_+$. Since J_2 anticommutes with J_1 it interchanges E and E^* and hence defines an isomorphism $E \xrightarrow{\sim} E^*$. Now $g(\cdot, J_2 \cdot)$ defines a symplectic form ω_E on E . Let $H = M \times \mathbb{C}^2 = M \times (\mathbb{C}h_1 \oplus \mathbb{C}h_2)$ be the trivial bundle with 2-form $\omega_H = h_1^* \wedge h_2^*$. Then we can identify

$$TM = E \oplus E^* = E \oplus E = h_1 \otimes E \oplus h_2 \otimes E = H \otimes E.$$

We check that under this identification we have $g = \omega_H \otimes \omega_E$. Note that both sides vanish on $h_1 \otimes E$ and $h_2 \otimes E$ and $\omega_H(h_1, h_2) = 1$. We calculate for $e, e' \in E = E_+ = h_1 \otimes E$:

$$g(e, J_2 e') = \omega_E(e, e') = \omega_H(h_1, h_2) \omega_E(e, e') = (\omega_H \otimes \omega_E)(h_1 \otimes e, h_2 \otimes e').$$

Hence we have a Grassmann structure. The eigenspaces $E_\pm$ of the parallel endomorphism J_1 are invariant under parallel transport. Therefore the Levi-Civita connection ∇ induces a connection ∇^E in the bundle E . Since $\nabla g = 0$ and $\nabla J_2 = 0$ we have $\nabla^E \omega_E = 0$. We define a flat connection ∇^H on the trivial bundle $H = M \times \mathbb{C}^2$ by the condition $\nabla^H h_1 = \nabla^H h_2 = 0$. Then $\nabla = \nabla^H \otimes \text{Id} + \text{Id} \otimes \nabla^E$. So the Grassmann structure is half-flat.

Finally, using the standard identification $\mathbb{C}^2 = \mathbb{H}$, one can easily check that the J_α are given by (4.1). $\square$

4.2 Complexification of real hyper-Kähler manifolds

Let (M, g, J_α) be a (real) hyper-Kähler manifold. We will assume that it is real analytic. This is automatically true if the metric g is positive definite since it is Ricci-flat and a fortiori Einstein. Using analytic continuation we can extend (M, g, J_α) to a complex hyper-Kähler manifold $(M^\mathbb{C}, g^\mathbb{C}, J_\alpha^\mathbb{C})$ equipped with an antiholomorphic involution T . In complex local coordinates $z^j = x^j + iy^j$ which are extension of real analytic coordinates x^j, y^j the involution is given by the complex conjugation $z^j \rightarrow \bar{z}^j = x^j - iy^j$. We can reconstruct the (real) hyper-Kähler manifold as the fixed point set of T . We will call (M, g, J_α) a **real form** of $(M^\mathbb{C}, g^\mathbb{C}, J_\alpha^\mathbb{C})$ and $(M^\mathbb{C}, g^\mathbb{C}, J_\alpha^\mathbb{C})$ the complexification of (M, g, J_α) .

In general a complex hyper-Kähler manifold has no real form. A necessary condition is that the holonomy group of ∇^E is contained in $\text{Sp}(k, l)$, $n = k + l$, and hence preserves a quaternionic

structure. Then we can define a parallel antilinear endomorphism field $j_E : E \rightarrow E$ such that $j_E^2 = -1$ and $\omega_E(j_E x, j_E y) = \overline{\omega_E(x, y)}$ for all $x, y \in E$, where the bar denotes complex conjugation. We define a parallel antilinear endomorphism field $j_H : H \rightarrow H$ as the left-multiplication by the quaternion j on $H = M \times \mathbb{H}$. Then $\rho = j_H \otimes j_E$ defines a field of real structures in $TM = H \otimes E$. We denote by $\mathcal{D} \subset TM$ the real eigenspace distribution of ρ with eigenvalue 1. Here TM is considered as real tangent bundle of the real manifold M . If $M^\rho \subset M$ is a leaf of $\mathcal{D}$ of real dimension $4n$ then the data (g, J_α) induce on M^ρ a (real) hyper-Kähler structure.

4.3 Complex hyper-Kähler symmetric spaces

A complex Riemannian symmetric space is a complex Riemannian manifold (M, g) such that any point is an isolated fixed point of an isometric holomorphic involution. Like in the real case one can prove that it admits a transitive complex Lie group of holomorphic isometries and that any simply connected complex Riemannian symmetric M is associated to a complex symmetric decomposition

$$\mathfrak{g} = \mathfrak{k} + \mathfrak{m}, \quad [\mathfrak{k}, \mathfrak{k}] \subset \mathfrak{k}, \quad [\mathfrak{k}, \mathfrak{m}] \subset \mathfrak{m}, \quad [\mathfrak{m}, \mathfrak{m}] = \mathfrak{k} \quad (4.2)$$

of a complex Lie algebra $\mathfrak{g}$ together with an $\text{ad}_{\mathfrak{k}}$ -invariant complex scalar product on $\mathfrak{m}$. More precisely $M = G/K$, where G is the simply connected complex Lie group with the Lie algebra $\mathfrak{g}$ and K is the (closed) connected subgroup associated with $\mathfrak{k}$. The holonomy group of such manifold is $H = \text{Ad}_K|_{\mathfrak{m}}$. Any pseudo-Riemannian symmetric space $M = G/K$ associated with a symmetric decomposition $\mathfrak{g} = \mathfrak{k} + \mathfrak{m}$ has a canonical complexification $M^{\mathbb{C}} = G^{\mathbb{C}}/K^{\mathbb{C}}$ defined by the complexification $\mathfrak{g}^{\mathbb{C}} = \mathfrak{k}^{\mathbb{C}} + \mathfrak{m}^{\mathbb{C}}$ of the symmetric decomposition. Proposition 1 remains true for complex Riemannian symmetric spaces. Ignoring the reality condition we obtain the following complex version of Theorem 1.

Theorem 2 *Let $S \in S^4 E$, $E = \mathbb{C}^{2n}$, be a quartic polynomial invariant under all endomorphisms $S_{e,e'} \in S^2 E = \text{sp}(E)$. Then it defines a complex hyper-Kähler symmetric space, which is associated with the following complex symmetric decomposition*

$$\mathfrak{g} = \mathfrak{h} + H \otimes E, \quad \mathfrak{h} = S_{E,E} = \text{span}\{S_{e,e'} | e, e' \in E\} \subset \text{sp}(E). \quad (4.3)$$

The bracket $\wedge^2(H \otimes E) \rightarrow \mathfrak{h}$ is given by (3.1). The complex hyper-Kähler symmetric space M associated to this symmetric decomposition is the quotient $M = M_S = G/K$, where G is the (complex) simply connected Lie group with Lie algebra $\mathfrak{g}$ and $K \subset G$ is the connected (and closed) subgroup with Lie algebra $\mathfrak{k} = \mathfrak{h}$.

Moreover any simply connected complex hyper-Kähler symmetric space can be obtained by this construction. Two complex hyper-Kähler symmetric spaces M_S and $M_{S'}$ defined by quartics

S and S' are isomorphic if and only if S and S' are in the same orbit of $\text{Aut}(E, \omega_E) = \text{Sp}(E) \cong \text{Sp}(n, \mathbb{C})$.

Corollary 1 *There is a natural bijection between simply connected complex hyper-Kähler symmetric spaces of dimension $4n$ up to isomorphism and $\text{Sp}(n, \mathbb{C})$ -orbits on the space of quartic polynomials $S \in S^4 E$ in the symplectic vector space $E = \mathbb{C}^{2n}$ such that*

$$S_{e,e'} \cdot S = 0 \quad \text{for all } e, e' \in E. \quad (4.4)$$

4.4 Classification of complex hyper-Kähler symmetric spaces

The following complex version of Proposition 4 (with similar proof) will be a crucial step in the classification of complex hyper-Kähler symmetric spaces.

Proposition 6 *Let $(M = G/K, g, J_\alpha)$ be a simply connected complex hyper-Kähler symmetric space. Then the holonomy group of M is solvable and M admits a transitive solvable Lie group of automorphisms.*

Due to Corollary 1 the classification of simply connected complex hyper-Kähler symmetric spaces reduces to the determination of quartic polynomials S satisfying (4.4). Below we will determine all such polynomials. We will prove that the following example gives all such polynomials.

Example 1: Let $E = E_+ \oplus E_-$ be a Lagrangian decomposition, i.e. $\omega(E_\pm, E_\pm) = 0$, of the symplectic vector space $E = \mathbb{C}^{2n}$. Then any polynomial $S \in S^4 E_+ \subset S^4 E$ satisfies the condition (4.4) and defines a simply connected complex hyper-Kähler symmetric space M_S with Abelian holonomy algebra $\mathfrak{h} = S_{E_+, E_+} \subset S^2 E_+ \subset S^2 E = \mathfrak{sp}(E)$.

In fact, since E_+ is Lagrangian the endomorphisms from $S^2 E_+$ form an Abelian subalgebra of $\mathfrak{sp}(E)$, which acts trivially on E_+ and hence on $S^4 E_+$.

Theorem 3 *Let $S \in S^4 E$ be a quartic polynomial satisfying (4.4). Then there exists a Lagrangian decomposition $E = E_+ \oplus E_-$ such that $S \in S^4 E_+$.*

Proof: According to Theorem 2 the quartic S defines a hyper-Kähler symmetric space with holonomy Lie algebra $\mathfrak{h} = S_{E,E}$. Since, by Proposition 6, $\mathfrak{h}$ is solvable, Lie's theorem implies the existence of a one-dimensional $\mathfrak{h}$ -invariant subspace $P = \mathbb{C}p \subset E$. There exists an ω -nondegenerate subspace $W \subset E$ such that the ω -orthogonal complement of P is $P^\perp = P \oplus W$.

We choose a vector $q \in E$ such that $\omega(p, q) = 1$ and $\omega(W, q) = 0$ and put $Q := \mathbb{C}q$. Then we have

$$E = P \oplus W \oplus Q.$$

Since $\mathfrak{h}$ preserves P we have the following inclusion

$$\mathfrak{h} \subset PE + W^2 = P^2 + PW + PQ + W^2,$$

where we use the notation $XY = X \vee Y$ for the symmetric product of subspaces $X, Y \subset E$. Then the second prolongation $\mathfrak{h}^{(2)} = \{T \in S^4 E | T_{e,e'} \in \mathfrak{h} \text{ for all } e, e' \in E\}$ has the following inclusion

$$\mathfrak{h}^{(2)} \subset P^3 E + P^2 W^2 + PW^3 + W^4 = P^4 + P^3 Q + P^3 W + P^2 W^2 + PW^3 + W^4. \quad (4.5)$$

Indeed $\mathfrak{h}^{(2)} \subset \mathfrak{h}^2 = P^4 + P^3 Q + P^3 W + P^2 Q^2 + P^2 WQ + P^2 W^2 + PQW^2 + PW^3 + W^4$. The projection $\mathfrak{h}^{(2)} \rightarrow P^2 Q^2 + P^2 WQ + PQW^2$ is zero because otherwise $S_{q,q} \in \mathfrak{h} \subset PE + W^2$ would have a nonzero projection to $Q^2 + WQ$ or $S_{w,q} \in \mathfrak{h}$ would have a nonzero projection to QW for appropriate choice of $w \in W$. By (4.5) we can write the quartic S as

$$S = p^3(\lambda p + \mu q + w_0) + p^2 B + pC + D,$$

where $\lambda, \mu \in \mathbb{C}$, $w_0 \in W$, $B \in S^2 W$, $C \in S^3 W$ and $D \in S^4 W$. From now on we will identify $S^d E$ with the space $\mathbb{C}[E^*]^{(d)}$ of homogeneous polynomials on E^* of degree d . Then the ω -contraction $T_x = \iota_{\omega x} T = T(\omega x, \dots)$ of a tensor $T \in S^d E$ with a vector $x \in E$ is identified with the following homogeneous polynomial of degree $d - 1$:

$$T_x = \frac{1}{d} \partial_{\omega x} T,$$

where $\partial_{\omega x} T$ is the derivative of the polynomial $T \in \mathbb{C}[E^*]^{(d)}$ in the direction of $\omega x = \omega(x, \cdot) \in E^*$. For example $p_q = \langle p, \omega q \rangle = \omega(q, p) = \partial_{\omega q} p = -\partial_{p^*} p = -1 = -q_p$.

From $S_{p,q} = -\frac{1}{4}\mu p^2$ and the condition $S_{p,q} \cdot S = 0$ we obtain $\mu = 0$, since $p^2 \cdot S = \mu p^4$. This implies $S_{p,\cdot} = 0$. Next we compute:

$$\begin{aligned} S_{q,q} &= \frac{1}{6}(6\lambda p^2 + 3pw_0 + B) \\ S_{q,w} &= -\frac{1}{12}(-3p^2\omega(w_0, w) + 2p\partial_{\omega w} B + \partial_{\omega w} C) \\ &= -\frac{1}{12}(-3p^2\omega(w_0, w) + 4pB_w + 3C_w) \\ S_{w,w'} &= \frac{1}{6}(p^2 B_{w,w'} + 3pC_{w,w'} + 6D_{w,w'}) \end{aligned}$$

for any $w, w' \in W$.

Now the condition (4.4) can be written as follows:

$$\begin{aligned}
0 &= 6S_{q,q} \cdot S = (3pw_0 + B) \cdot S = \left(\frac{3}{2}(p \otimes w_0 + w_0 \otimes p) + B\right) \cdot S \\
&= \frac{3}{2}(2p^3Bw_0 + 3p^2Cw_0 + 4pDw_0) + p^3Bw_0 + p^2B \cdot B + pB \cdot C + B \cdot D \\
&= -2p^3Bw_0 + \frac{9}{2}p^2Cw_0 + p(6Dw_0 + B \cdot C) + B \cdot D.
\end{aligned}$$

Note that $Bw_0 = -B_{w_0}$ and $B \cdot B = [B, B] = 0$.

$$\begin{aligned}
0 &= -12S_{q,w} \cdot S = (4pB_w + 3C_w) \cdot S \\
&= 2(p^4\omega(B_w, w_0) + 2p^3B^2w - 3p^2C_{Bw} - 4pD_{Bw}) \\
&\quad + 3(p^3C_w w_0 + p^2C_w \cdot B + pC_w \cdot C + C_w \cdot D) \\
&= 2p^4\omega(B_w, w_0) + p^3(4B^2w + 3C_w w_0) + p^2(-6C_{Bw} + 3C_w \cdot B) \\
&\quad + p(-8D_{Bw} + 3C_w \cdot C) + 3C_w \cdot D \\
0 &= 2S_{w,w'} \cdot S = \frac{1}{2}(p \otimes C_{w,w'} + C_{w,w'} \otimes p) \cdot S + 2D_{w,w'} \cdot S \\
&= \frac{1}{2}(p^4\omega(C_{w,w'}, w_0) - 2p^3BC_{w,w'} + 3p^2C_{C_{w,w'}} + 4pD_{C_{w,w'}}) + \\
&\quad 2(p^3D_{w,w'}w_0 + p^2D_{w,w'} \cdot B + pD_{w,w'} \cdot C + D_{w,w'} \cdot D) \\
&= \frac{1}{2}p^4\omega(C_{w,w'}, w_0) + p^3(-BC_{w,w'} + 2D_{w,w'}w_0) + \\
&\quad p^2\left(\frac{3}{2}C_{C_{w,w'}} + 2D_{w,w'} \cdot B\right) + p(2D_{C_{w,w'}} + 2D_{w,w'} \cdot C) + 2D_{w,w'} \cdot D
\end{aligned}$$

This gives the following system of equations:

- (1) $Bw_0 = 0$
- (2) $C_{w_0} = 0$
- (3) $6Dw_0 + B \cdot C = 0$
- (4) $B \cdot D = 0$
- (5) $\omega(B_w, w_0) = 0$
- (6) $4B^2w + 3C_w w_0 = 0$
- (7) $-2C_{Bw} + C_w \cdot B = 0$
- (8) $-8D_{Bw} + 3C_w \cdot C = 0$
- (9) $C_w \cdot D = 0$
- (10) $\omega(C_w w', w_0) = 0$
- (11) $-BC_w w' + 2D_{w,w'}w_0 = 0$
- (12) $\frac{3}{2}C_{C_w w'} + 2D_{w,w'} \cdot B = 0$
- (13) $D_{C_w w'} + D_{w,w'} \cdot C = 0$
- (14) $D_{w,w'} \cdot D = 0$

Note that (5) and (10) follow from (1) and (2) and that using (2) equation (6) says that the endomorphism B has zero square:

$$(6') \quad B^2 = 0.$$

Eliminating D_{w_0} in equations (3) and (11) we obtain:

$$(15) \quad 0 = (B \cdot C)_w w' + 3BC_w w' = BC_w w' - C_{Bw} w' - C_w Bw' + 3BC_w w' = 4BC_w w' - C_{Bw} w' - C_w Bw'.$$

We can rewrite (7) as:

$$(7') \quad -2C_{Bw} w' + C_w Bw' - BC_w w' = 0.$$

Eliminating $C_{Bw} w'$ in (7') and (15) we obtain:

$$(16) \quad -3BC_w w' + C_w Bw' = 0.$$

Since the first summand is symmetric in w and w' we get

$$(17) \quad C_w Bw' = C_{w'} Bw = C_{Bw} w'.$$

Now using (17) we can rewrite (15) as:

$$(15') \quad 2BC_w w' - C_w Bw' = 0.$$

The equations (15') and (16) show that $BC_w w' = C_w Bw' = C_{Bw} w' = 0$ and hence also $B \cdot C = 0$.

This implies $D_{w_0} = 0$, by (3). Now we can rewrite (1-14) as:

$$Bw_0 = C_{w_0} = D_{w_0} = 0 \tag{4.6}$$

$$B \cdot C = B \cdot D = 0 \tag{4.7}$$

$$B^2 = 0 \tag{4.8}$$

$$C_{Bw} = C_w B = BC_w = 0 \tag{4.9}$$

$$-8D_{Bw} + 3C_w \cdot C = 0 \tag{4.10}$$

$$C_w \cdot D = 0 \tag{4.11}$$

$$\frac{3}{2}C_{C_w w'} + 2D_{w, w'} \cdot B = 0 \tag{4.12}$$

$$D_{C_w w'} + D_{w, w'} \cdot C = 0 \tag{4.13}$$

$$D_{w, w'} \cdot D = 0 \tag{4.14}$$

Now to proceed further we decompose $K := \ker B = W_0 \oplus W'$, where $W_0 = \ker \omega|_K$ and W' is a (nondegenerate) complement. Let us denote by W_1 a complement to K in W such that $\omega(W', W_1) = 0$. Then $W_0 + W_1$ is the ω -orthogonal complement to the B -invariant nondegenerate subspace W' . This shows that $BW_1 \subset (W_0 + W_1) \cap K = W_0$. Moreover since $W_1 \cap K = 0$ the map $B : W_1 \rightarrow W_0$ is injective and hence $\dim W_1 \leq \dim W_0$. On the other hand $\dim W_1 \geq \dim W_0$, since W_0 is an isotropic subspace of the symplectic vector space $W_0 + W_1$. This shows that $B : W_1 \rightarrow W_0$ is an isomorphism.

Lemma 3 $C \in S^3K$ and $D \in S^4K$.

Proof: Since $W_0 = BW$ the equation (4.9) shows that $C_{W_0} = 0$, which proves the first statement. From (4.10) and the identity

$$(C_x \cdot C)_y = [C_x, C_y] - C_{C_{xy}} \quad (4.15)$$

we obtain

$$D_{Bx,y} + D_{By,x} = \frac{3}{8}((C_x \cdot C)_y + (C_y \cdot C)_x) = -\frac{3}{4}C_{C_{xy}}. \quad (4.16)$$

The equation $B \cdot D = 0$ (4.7) reads:

$$0 = (B \cdot D)_{x,y} = [B, D_{x,y}] - D_{Bx,y} - D_{x,By}.$$

Using this (4.12) yields:

$$D_{Bx,y} + D_{By,x} = [B, D_{x,y}] = -D_{x,y} \cdot B = \frac{3}{4}C_{C_{xy}}. \quad (4.17)$$

Now from (4.16) and (4.17) we obtain that

$$0 = C_{C_{xy}}z = C_zC_{xy}$$

for all $x, y, z \in W$. This implies $[C_x, C_y] = 0$ for all $x, y \in W$ and hence

$$C_x \cdot C = 0 \quad (4.18)$$

for all $x \in W$, by (4.15). Finally this shows that $D_{W_0} = 0$ by (4.10). This proves the second statement. $\square$

Lemma 4 $D_{x,y}C_z = C_zD_{x,y} = 0$ for all $x, y, z \in W$.

Proof: Using (4.13) we compute:

$$D_{x,y}C_zw = D_{C_{zw},xy} = -(D_{z,w} \cdot C)_{xy} = -([D_{z,w}, C_x]y - C_{D_{z,w}xy}). \quad (4.19)$$

From (4.11) we get:

$$0 = (C_x \cdot D)_{z,wy} = [C_x, D_{z,w}]y - D_{C_{xz},wy} - D_{z,C_{xy}} = C_xD_{z,w}y - D_{z,w}C_{xy} - D_{y,w}C_{xz} - D_{z,y}C_{xw},$$

and hence:

$$[D_{z,w}, C_x]y = -D_{y,w}C_{xz} - D_{z,y}C_{xw},$$

and

$$C_{D_{z,w}xy} = C_yD_{z,w}x = D_{z,w}C_{xy} + D_{x,w}C_yz + D_{z,x}C_yw.$$

Now we eliminate the CD -terms from (4.19) arriving at:

$$D_{x,y}C_z w = (D_{y,w}C_x z + D_{z,y}C_x w + D_{z,w}C_x y + D_{x,w}C_y z + D_{z,x}C_y w). \quad (4.20)$$

Considering all the permutations of (x, y, z, w) we get 6 homogeneous linear equations for the 6 terms of equation (4.20) with the matrix:

$$\begin{pmatrix} -1 & 1 & 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & 1 & 1 & 1 \\ 1 & 1 & 1 & -1 & 1 & 1 \\ 1 & 1 & 1 & 1 & -1 & 1 \\ 1 & 1 & 1 & 1 & 1 & -1 \end{pmatrix}$$

This is the matrix of the endomorphism $-2 \text{Id} + e \otimes e$ in the arithmetic space $\mathbb{R}^6$, where $e = e_1 + \dots + e_6$; (e_i) the standard basis. It has eigenvalues $(4, -2, -2, -2, -2, -2)$. This shows that the matrix is nondegenerate and proves the lemma. $\square$

For a symmetric tensor $T \in S^d W$ we denote by

$$\Sigma_T := \text{span}\{T_{x_1, x_2, \dots, x_{d-2}} x_{d-1} | x_1, x_2, \dots, x_{d-1} \in W\} \subset W$$

the **support** of T .

Lemma 5 *The supports of the tensors $B \in S^2 W$, $C \in S^3 W$ and $D \in S^4 W$ admit the following inclusions*

$$\Sigma_B + \Sigma_C \subset \ker B \cap \ker C \cap \ker D, \quad \Sigma_D \subset \ker B \cap \ker C.$$

Moreover $\Sigma_B + \Sigma_C$ is isotropic and $\omega(\Sigma_D, \Sigma_B + \Sigma_C) = 0$.

Proof: The first statement follows from $B^2 = BC_x = BD_{x,y} = C_x B = C_x C_y = C_x D_{y,z} = D_{x,y} B = D_{x,y} C_z = 0$ for all $x, y, z \in W$. The second statement follows from the first and the definition of support, e.g. if $z = C_x y \in \Sigma_C$ and $w \in \Sigma_B + \Sigma_C + \Sigma_D \subset \ker C$ we compute:

$$\omega(z, w) = \omega(C_x y, w) = -\omega(y, C_x w) = 0.$$

$\square$

Lemma 6 *The Lie algebra $D_{W,W} \subset S^2 W \cong \text{sp}(W)$ is solvable.*

Proof: This follows from Proposition 6, since $D \in S^4 W$ satisfies (4.4) and hence defines a complex hyper-Kähler symmetric space with holonomy Lie algebra $D_{W,W}$. It also follows from

the solvability of $S_{E,E}$ as we show now. In terms of the decomposition $E = P + W + Q$ an endomorphism

$$S_{x,y} = (\lambda p^4 + p^3 w_0 + p^2 B + pC + D)_{x,y} = (p^2 B + pC + D)_{x,y} = B(x,y)p^2 - pC_{xy} + D_{x,y},$$

where $x, y \in W$, is represented by

$$\begin{pmatrix} 0 & -(C_x y)^t & B(x,y) \\ 0 & D_{x,y} & -C_{xy} \\ 0 & 0 & 0 \end{pmatrix}.$$

Since the Lie algebra $S_{E,E}$ is solvable this implies that the Lie algebra $D_{W,W}$, which corresponds to the induced representation of $S_{E,E}$ on $P^\perp/P \cong W$, is also solvable. $\square$

Lemma 7

$$\omega(w_0, \Sigma_B + \Sigma_C + \Sigma_D) = 0.$$

Proof: Note that $w_0 \in \ker B \cap \ker C \cap \ker D$, due to equations (1-3) and (4.7). This implies the lemma. In fact, if e.g. $y = Bx \in \Sigma_B$ then

$$\omega(w_0, y) = \omega(w_0, Bx) = -\omega(Bw_0, x) = 0,$$

which shows that $\omega(w_0, \Sigma_B) = 0$ $\square$

Now to finish the proof of Theorem 3 we will use induction on the dimension $\dim E = 2n$. If $n = 1$ the (solvable) holonomy algebra $\mathfrak{h}$ is a proper subalgebra of $S^2 E \cong \mathfrak{sl}(2, \mathbb{C})$. Without loss of generality we may assume that either

- a) $\mathfrak{h} = \mathbb{C}p^2$ or
- b) $\mathfrak{h} = \mathbb{C}pq$ or
- c) $\mathfrak{h} = \mathbb{C}p^2 + \mathbb{C}pq$,

where (p, q) is a symplectic basis of E . In the all three cases the Lie algebra $S_{E,E} = \mathfrak{h} \subset \mathbb{C}p^2 + \mathbb{C}pq$ and hence

$$S = \lambda p^4 + \mu p^3 q.$$

In the cases b) and c) we have that $pq \in \mathfrak{h}$ and since

$$pq \cdot S = \frac{1}{2}(p \otimes q + q \otimes p) \cdot S = -2\lambda p^4 - \mu p^3 q = 0$$

it follows that $S = 0$. In the case a) from $S_{E,E} = \mathfrak{h} = \mathbb{C}p^2$ we have that $S = \lambda p^4$. This tensor is invariant under $\mathfrak{h} = \mathbb{C}p^2$ and belongs to the fourth symmetric power of the Lagrangian subspace $\mathbb{C}p \subset E$. This establishes the first step of the induction. Now by induction using equation (4.14) and Lemma 6 we may assume that Σ_D is isotropic. Now Lemma 5 and Lemma 7 show that

$\mathbb{C}w_0 + \Sigma_B + \Sigma_C + \Sigma_D$ is isotropic and hence is contained in some Lagrangian subspace $E_+ \subset E$. This implies that $S \in S^4E_+$. $\square$

Now we give a necessary and sufficient condition for a symmetric manifold $M = M_S$, $S \in S^4E_+$, to have no flat de Rham factor.

Proposition 7 *The complex hyper-Kähler symmetric space M_S , $S \in S^4E_+$, has no flat de Rham factor if and only if the support $\Sigma_S = E_+$.*

Proof: If $M = M_S = G/K$ has a flat factor M_0 , such that $M = M_1 \times M_0$, then this induces a decomposition $E = E^1 \oplus E^0$ and $S \in S^4E_1$; hence $\Sigma_S \subset E_1 \cap E_+ \neq E_+$. Conversely let $S \in S^4E_+$, assume that $E_+^1 = \Sigma_S \subset E_+$ is a proper subspace and choose a complementary subspace E_+^0 . We denote by E_-^1 and E_-^0 the annihilator of ωE_+^0 and ωE_+^1 respectively. Let us denote $E^1 = E_+^1 \oplus E_-^1$, $E^0 = E_+^0 \oplus E_-^0$, $\mathfrak{m}^1 = H \otimes E^1$ and $\mathfrak{m}^0 = H \otimes E^0$. Then $E^0, E^1 \subset E$ are ω -nondegenerate complementary subspaces and $\mathfrak{m}^0, \mathfrak{m}^1 \subset \mathfrak{m} = T_oM$ are g -nondegenerate complementary subspaces. Since $S \in S^4E_+^1$ the Lie algebra $\mathfrak{g} = \mathfrak{h} + \mathfrak{m} = (\mathfrak{h} + \mathfrak{m}^1) \oplus \mathfrak{m}^0$ has the Abelian direct summand $\mathfrak{m}^0$, see (3.1), which gives rise to a flat factor $M^0 \subset M_S = M^1 \times M^0$.

Theorem 4 *Any simply connected complex hyper-Kähler symmetric space without flat de Rham factor is isomorphic to a complex hyper-Kähler symmetric space of the form M_S , where $S \in S^4E_+$ and $E_+ \subset E$ is a Lagrangian subspace of the complex symplectic vector space $E = \mathbb{C}^{2n}$. Moreover there is a natural 1-1 correspondence between simply connected complex hyper-Kähler symmetric spaces without flat factor up to isomorphism and orbits $\mathcal{O}$ of the group $\text{Aut}(E, \omega, E_+) = \{A \in \text{Sp}(E) | AE_+ = E_+\} \cong \text{GL}(E_+) \cong \text{GL}(n, \mathbb{C})$ on the space S^4E_+ such that $\Sigma_S = E_+$ for all $S \in \mathcal{O}$.*

Proof: This is a corollary of Theorem 2, Theorem 3 and Proposition 7. $\square$

Let $M = G/K$ be a simply connected complex hyper-Kähler symmetric space without flat factor. By Theorem 3 and Proposition 7 it is associated to quartic polynomial $S \in S^4E_+$ with support $\Sigma_S = E_+$. Now we describe the Lie algebra $\text{aut}(M_S)$ of the full group of automorphisms, i.e. isometries which preserve the hypercomplex structure, of M_S .

Theorem 5 *Let $M_S = G/K$ be as above. Then the full automorphism algebra is given by*

$$\text{aut}(M_S) = \text{aut}(S) + \mathfrak{g},$$

where $A \in \text{aut}(S) = \{B \in \text{gl}(E_+) | B \cdot S = 0\}$ acts on $\mathfrak{g} = \mathfrak{h} + \mathfrak{m}$ as follows. It preserves the decomposition and acts on $\mathfrak{h} = S_{E,E}$ by

$$[A, S_{x,y}] = S_{Ax,y} + S_{x,Ay}$$

for all $x, y \in E$ and on $\mathfrak{m} = H \otimes E$ by

$$[A, h \otimes e] = h \otimes Ae,$$

where $\mathfrak{gl}(E_+)$ is canonically embedded into $\mathfrak{sp}(E)$.

Proof: By (the complex version of) Proposition 1 it is sufficient to determine the centralizer $\mathfrak{c}$ of $\mathfrak{sp}(1, \mathbb{C})$ in the full isotropy algebra $\tilde{\mathfrak{h}} = \mathfrak{aut}(R) \supset \mathfrak{sp}(1, \mathbb{C}) \oplus \mathfrak{h}$. Equation (2.2) shows that

$$\mathfrak{c} = \{\text{Id} \otimes A \mid A \in \mathfrak{sp}(E), A \cdot S = [A, S(\cdot, \cdot)] - S(A \cdot, \cdot) - S(\cdot, A \cdot) = 0\}.$$

From $A \cdot S = 0$ we obtain that the commutator $[A, S_{x,y}] = S_{Ax,y} + S_{x,Ay}$ for all $x, y \in E$ and $A\Sigma_S = AE_+ \subset E_+$. This implies $\mathfrak{c} = \mathfrak{aut}(S)$. $\square$

5 Classification of hyper-Kähler symmetric spaces

Using the description of complex hyper-Kähler symmetric spaces given in Theorem 4 we will now classify (real) hyper-Kähler symmetric spaces. Recall that a simply connected pseudo-Riemannian manifold is called **indecomposable** if it is not a Riemannian product of two pseudo-Riemannian manifolds. Any simply connected pseudo-Riemannian manifold can be decomposed into the Riemannian product of indecomposable pseudo-Riemannian manifolds. By Wu's theorem [W] a simply connected pseudo-Riemannian manifold is indecomposable if and only if its holonomy group is **weakly irreducible**, i.e. has no invariant proper nondegenerate subspaces. Therefore it is sufficient to classify (real) hyper-Kähler symmetric spaces with *indecomposable* holonomy.

Let $(M = G/K, g, J_\alpha)$ be a hyper-Kähler symmetric space associated to a symmetric decomposition (2.1). The complexified tangent space of M is identified with $\mathfrak{m}^\mathbb{C} = H \otimes E$, the tensor product of two complex symplectic vector spaces with quaternionic structure j_H and j_E such that $\rho = j_H \otimes j_E$ is the complex conjugation of $\mathfrak{m}^\mathbb{C}$ with respect to $\mathfrak{m}$. By Theorem 1 it is defined by a quartic polynomial $S \in S^4 E$ satisfying the conditions of the theorem. Moreover the holonomy algebra $\mathfrak{h}$ acts trivially on H and is identified with the real form of the complex Lie algebra $S_{E,E} \subset \mathfrak{sp}(E)$ given by $\mathfrak{h} = \text{span}\{S_{je,e'} - S_{e,je'} \mid e, e' \in E\} = \{A \in S_{E,E} \mid [A, j_E] = 0\} \subset \mathfrak{sp}(E)^{j_E}$.

The quartic polynomial S defines also a complex hyper-Kähler symmetric space $M^\mathbb{C} = G^\mathbb{C}/K^\mathbb{C}$, which is the complexification of $M = G/K$. By Theorem 3, $S \in S^4 L$ for some Lagrangian subspace $L \subset E$. Recall that the symplectic form $\omega = \omega_E$ together with the quaternionic structure $j = j_E$ define a Hermitian metric $\gamma = \gamma_E = \omega_E(\cdot, j_E \cdot)$ of (real) signature $(4k, 4l)$, $n = k + l$, which coincides with the signature of the pseudo-Riemannian metric g (we normalize $\gamma_H = \omega_H(\cdot, j_H \cdot)$ to be positive definite). We may decompose γ -orthogonally

$L = L^0 \oplus L^+ \oplus L^-$, such that γ vanishes on L^0 is positive definite on L^+ and negative definite on L^- .

Lemma 8 (i) $jL^0 = L^0$ and

(ii) $L^+ + L^- + jL^+ + jL^- \subset E$ is an ω -nondegenerate and γ -nondegenerate $\mathfrak{h}$ -invariant subspace (with trivial action of $\mathfrak{h}$).

Proof: We show first that $L + jL^0$ is ω -isotropic and hence $L + jL^0 = L$ since L is Lagrangian. Indeed $L \supset L^0$ is ω -isotropic and also jL^0 because ω is j -invariant. So it suffices to remark that $\omega(L, jL^0) = 0$:

$$\omega(L, jL^0) = \gamma(L, L^0) = 0.$$

This implies that $jL^0 \subset L$. Since $\gamma(L, jL^0) = -\omega(L, L^0) = 0$, we conclude that $jL^0 \subset \ker \gamma|_L = L^0$. This proves (i).

To prove (ii) it is sufficient to check that the subspace $L^+ + L^- + jL^+ + jL^- \subset E$ is nondegenerate with respect to γ , since it is j -invariant. First we remark that γ is positive definite on L^+ and jL^+ and negative definite on L^- and jL^- , due to the j -invariance of γ : $\gamma(jx, jx) = \gamma(x, x)$, $x \in E$. So to prove (ii) it is sufficient to check that $jL^+ \oplus jL^-$ is γ -orthogonal to the γ -nondegenerate vector space $L^+ + L^-$:

$$\gamma(L^+ + L^-, jL^+ + jL^-) = \omega(L^+ + L^-, L^+ + L^-) = 0.$$

□

By Theorem 1 the quartic polynomial S must satisfy the reality condition $[S_{je,e'} - S_{e,je'}, j] = 0$. Now we describe all such polynomials.

The quaternionic structure j on E is **compatible** with ω , i.e. $\omega(jx, jy) = \overline{\omega(x, y)}$ for all $x, y \in E$ and it induces a real structure (i.e. an antilinear involution) $\tau := j \otimes j \otimes \cdots \otimes j$ on all even powers $S^{2r}E \subset E \otimes E \otimes \cdots \otimes E$. For $S \in S^{2r}E$ and $x_1, \dots, x_{2r} \in E$ we have

$$(\tau S)(x_1, \dots, x_{2r}) = \overline{S(jx_1, \dots, jx_{2r})}.$$

Note that the fixed point set $\text{sp}(E)^\tau = \{A \in \text{sp}(E) | [A, j] = 0\} \cong \text{sp}(k, l)$.

Proposition 8 Let (E, ω, j) be a complex symplectic vector space with a quaternionic structure j such that $\omega(jx, jy) = \overline{\omega(x, y)}$ for all $x, y \in E$. Then a quartic polynomial $S \in S^4E$ satisfies the reality condition $[S_{je,e'} - S_{e,je'}, j] = 0$ if and only if $S \in (S^4E)^\tau = \text{span}\{T + \tau T | T \in S^4E\}$.

Proof: The reality condition for $S \in S^4E$ can be written as

$$[S_{jx, jy} + S_{x, y}, j]z = 0$$

for all $x, y, z \in E$. Contracting this vector equation with $ju \in E$ by means of ω and using the compatibility between j and ω we obtain the equivalent condition

$$\begin{aligned} 0 &= -\omega(jw, [S_{jx, jy} + S_{x, y}, j]z) \\ &= S(jx, jy, jz, jw) - \overline{S(jx, jy, z, w)} + S(x, y, jz, jw) - \overline{S(x, y, z, w)}. \end{aligned} \quad (5.1)$$

Now putting $x = y = z = w = u$ we obtain:

$$0 = S(ju, ju, ju, ju) - \overline{S(ju, ju, u, u)} + S(u, u, ju, ju) - \overline{S(u, u, u, u)}$$

and putting $x = iu$ and $y = z = w = u$ we obtain:

$$0 = -iS(ju, ju, ju, ju) - \overline{iS(ju, ju, u, u)} + iS(u, u, ju, ju) + \overline{iS(u, u, u, u)}.$$

Comparing these two equations we get $S(ju, ju, ju, u) = \overline{S(u, u, u, u)}$, i.e. $S = \tau S$. This shows that the reality condition implies that $S \in (S^4 E)^\tau$. Conversely the condition $S = \tau S$ can be written as

$$S(jx, jy, jz, jw) = \overline{S(x, y, z, w)} \quad \text{for all } x, y, z, w \in E.$$

Changing $z \rightarrow jz$ and $w \rightarrow jw$ in this equation we obtain

$$S(jx, jy, z, w) = \overline{S(x, y, jz, jw)} \quad \text{for all } x, y, z, w \in E.$$

These two equations imply (5.1) and hence the reality condition. $\square$

Now we are ready to classify simply connected hyper-Kähler symmetric spaces. We will show that the following construction gives all such symmetric spaces.

Let (E, ω, j) be a complex symplectic vector space of dimension $2n$ with a quaternionic structure j such that $\omega(jx, jy) = \overline{\omega(x, y)}$ for all $x, y \in E$ and $E = E_+ \oplus E_-$ a j -invariant Lagrangian decomposition. Such a decomposition exists if and only if the Hermitian form $\gamma = \omega(\cdot, j\cdot)$ has real signature $(4m, 4m)$, where $\dim_{\mathbb{C}} E = 2n = 4m$. Then any polynomial $S \in (S^4 E_+)^\tau = S^4 E_+ \cap (S^4 E)^\tau$ satisfies the condition (4.4) and the reality condition, by Proposition 8. Hence by Theorem 1 it defines a (real) simply connected hyper-Kähler symmetric space M_S with Abelian holonomy algebra $\mathfrak{h} = (S_{E_+, E_+})^\tau = S_{E_+, E_+} \cap (S^2 E)^\tau = \text{span}\{S_{je, e'} - S_{e, je'} | e, e' \in E\} \subset \text{sp}(E)^\tau \cong \text{sp}(m, m)$.

Theorem 6 *Any simply connected hyper-Kähler symmetric space without flat de Rham factor is isomorphic to a hyper-Kähler symmetric space of the form M_S , where $S = T + \tau T$, $T \in S^4 E_+$ and $E_+ \subset E$ is a j -invariant Lagrangian subspace of the complex symplectic vector space E with compatible quaternionic structure j . A hyper-Kähler symmetric space of the form M_S has no flat factor if and only if its complexification has no flat factor, which happens if and only if the*

support $\Sigma_S = E_+$. Moreover there is a natural 1-1 correspondence between simply connected hyper-Kähler symmetric spaces without flat factor up to isomorphism and orbits $\mathcal{O}$ of the group $\text{Aut}(E, \omega, j, E_+) = \{A \in \text{Sp}(E) \mid [A, j] = 0, AE_+ = E_+\} \cong \text{GL}(m, \mathbb{H})$ on the space $(S^4 E_+)^{\tau}$ such that $\Sigma_S = E_+$ for all $S \in \mathcal{O}$.

Proof: Let M be a simply connected hyper-Kähler symmetric space. We first assume that it is indecomposable. Then the holonomy algebra $\mathfrak{h}$ is weakly irreducible. By Theorem 1, $M = M_S$ for some quartic polynomial $S \in S^4 E$ satisfying (4.4) and the reality condition (3.2). By Proposition 8 the reality condition means that $S \in (S^4 E)^{\tau}$. On the other hand, by Theorem 3 $S \in S^4 L$ for some Lagrangian subspace L of E . Now the weak irreducibility of $\mathfrak{h}$ and Lemma 8 imply that $L = L^0$ is j -invariant. This proves that $S \in (S^4 E_+)^{\tau}$, where $E_+ = L = L^0$ is a j -invariant Lagrangian subspace of E . This shows that M is obtained from the above construction. Any simply connected hyper-Kähler symmetric space M without flat factor is the Riemannian product of indecomposable ones, say $M = M_1 \times M_2 \times \cdots \times M_r$, and we may assume that $M_i = M_{S_i}$, $S_i \in S^4 E_i$. Therefore M is associated to the quartic polynomial $S = S_1 \oplus S_2 \oplus \cdots \oplus S_r \in S^4 E$, $E = E_1 \oplus E_2 \oplus \cdots \oplus E_r$. Moreover S satisfies (4.4) and (3.2) if the S_i satisfy (4.4) and (3.2). This shows that any simply connected hyper-Kähler symmetric space is obtained from the above construction.

It is clear that the complexification $M_S^{\mathbb{C}}$ has a flat factor if M_S has a flat factor. Conversely let us assume that $M_S^{\mathbb{C}}$ has a flat factor, hence $\Sigma_S \subset E_+$ is a proper subspace. Since $j\Sigma_S = jS_{E,E}E = S_{E,E}jE = S_{E,E}E = \Sigma_S$ there exists a j -invariant complementary subspace E'_+ in E_+ . Denote by E'_- the annihilator of Σ_S in E_- then $E' = E'_+ \oplus E'_-$ is an ω -nondegenerate and j -invariant subspace of E on which the holonomy $\mathfrak{h}^{\mathbb{C}} = S_{E,E} \subset S^2 \Sigma_S$ acts trivially. Then the corresponding real subspace $(H \otimes E')^{\rho} \subset \mathfrak{m} = (H \otimes E)^{\rho}$ is a g -nondegenerate subspace on which the holonomy $\mathfrak{h}$ acts trivially. By Wu's theorem [W] it defines a flat de Rham factor.

Now the last statement follows from the corresponding statement in Theorem 1. $\square$

Corollary 2 *Any hyper-Kähler symmetric space without flat factor has signature $(4m, 4m)$. In particular its dimension is divisible by 8.*

Corollary 3 *Let $M = M_S$ be a complex hyper-Kähler symmetric space without flat factor associated with a quartic $S \in S^4 E_+$, where $E_+ \subset E$ is a Lagrangian subspace. It admits a real form if and only if there exists a quaternionic structure j on E compatible with ω preserving E_+ such that $\tau S = S$, where τ is the real structure on $S^4 E$ induced by j . In particular $\dim_{\mathbb{C}} M$ has to be divisible by 8.*

Let $M = G/K$ be a simply connected hyper-Kähler symmetric space without flat factor. By Theorem 6 it is associated to a quartic polynomial $S \in (S^4 E_+)^{\tau}$ with support $\Sigma_S = E_+$. Now we describe the Lie algebra $\text{aut}(M_S)$ of the full group of automorphisms, i.e. isometries which preserve the hypercomplex structure of M_S .

Theorem 7 *Let $M_S = G/K$ be as above. Then the full automorphism algebra is given by*

$$\text{aut}(M_S) = \text{aut}(S) + \mathfrak{g},$$

where $\text{aut}(S) = \{A \in \mathfrak{gl}(E_+) \mid [A, j] = 0, A \cdot S = 0\}$ acts on

$$\mathfrak{g} = \mathfrak{h} + \mathfrak{m}, \quad \mathfrak{h} = \{A \in S_{E,E} \mid [A, j] = 0\} = \text{span}\{S_{jx,y} - S_{x,jy} \mid x, y \in E\}, \mathfrak{m} = (H \otimes E)^{\rho}$$

as in Theorem 5.

Proof: The proof is similar to that of Theorem 5. □

6 Low dimensional hyper-Kähler symmetric spaces

6.1 Complex hyper-Kähler symmetric spaces of dimension ≤ 8

Dimension 4

Assume that M is a simply connected complex hyper-Kähler symmetric space of dimension 4. Applying Theorem 4 we conclude that $M = M_S$ for some $S \in S^4 E_+$, where $E_+ \subset E$ is a one-dimensional subspace $E_+ = \mathbb{C}e$. This proves:

Theorem 8 *There exists up to isomorphism only one non-flat simply connected complex hyper-Kähler symmetric space of dimension 4: $M = M_S$ associated with the quartic $S = e^4$.*

Dimension 8

Any eight-dimensional simply connected complex hyper-Kähler symmetric space is associated with a quartic $S \in S^4 E_+$, where $E_+ \subset E$ is a Lagrangian subspace of $E = \mathbb{C}^4$. We denote by (e, e') a basis of E_+ .

Theorem 9 *Eight-dimensional simply connected complex hyper-Kähler symmetric space are in natural 1-1 correspondence with the orbits of the group $\text{CO}(3, \mathbb{C}) = \mathbb{C}^* \cdot \text{SO}(3, \mathbb{C})$ on the space $S_0^2 \mathbb{C}^3$ of traceless symmetric matrices. The complex hyper-Kähler symmetric space associated with a traceless symmetric matrix A is the manifold $M_{S(A)}$, where $S(A) \in S^4 \mathbb{C}^2$ is the quartic polynomial which corresponds to A under the $\text{SO}(3, \mathbb{C})$ -equivariant isomorphism $S_0^2 \mathbb{C}^3 \cong S_0^2 \wedge^2 \mathbb{C}^3 \cong S_0^2 S^2 \mathbb{C}^2 = S^4 \mathbb{C}^2$.*

The classification of $\mathrm{SO}(3, \mathbb{C})$ -orbits on $S_0^2\mathbb{C}^3$ was given by Petrov [P] in his classification of Weyl tensors of Lorentzian 4-manifolds.

Proof: By Theorem 4 the classification of eight-dimensional simply connected complex hyper-Kähler symmetric spaces reduces to the description of orbits of the group $\mathrm{GL}(E_+) = \mathrm{GL}(2, \mathbb{C})$ on $S^4\mathbb{C}^2 \subset S^2S^2\mathbb{C}^2$. Fixing a volume form σ on $\mathbb{C}^2$ we can identify $S^2\mathbb{C}^2$ with $\mathfrak{sp}(1, \mathbb{C}) \cong \mathfrak{so}(3, \mathbb{C})$. Then the Killing form B is an $\mathrm{SL}(2, \mathbb{C})$ -invariant and we have the $\mathrm{GL}(2, \mathbb{C})$ -invariant decomposition: $S^2S^2\mathbb{C}^2 = S_0^2S^2\mathbb{C}^2 \oplus \mathbb{C}B$. The action of $\mathrm{SL}(2, \mathbb{C})$ on $S^2\mathbb{C}^2$ is effectively equivalent to the adjoint action of $\mathrm{SO}(3, \mathbb{C})$. The problem thus reduces essentially to the determination of the orbits of $\mathrm{SO}(3, \mathbb{C})$ on $S_0^2\mathbb{C}^3$. $\square$

6.2 Hyper-Kähler symmetric spaces of dimension ≤ 8

By Corollary 3 the minimal dimension of non-flat hyper-Kähler symmetric spaces is 8.

Theorem 10 *Eight-dimensional simply connected hyper-Kähler symmetric space are in natural 1-1 correspondence with the orbits of the group $\mathbb{R}^+ \cdot \mathrm{SO}(3)$ on the space $S_0^2\mathbb{R}^3$ of traceless symmetric matrices. The hyper-Kähler symmetric space associated with a traceless symmetric matrix A is the manifold $M_{S(A)}$, where $S(A) \in (S^4\mathbb{C}^2)^\tau$ is the quartic polynomial which corresponds to A under the $\mathrm{SO}(3)$ -equivariant isomorphism $S_0^2\mathbb{R}^3 \cong (S^4\mathbb{C}^2)^\tau$.*

References

- [BD] M.L. Barberis, I. Dotti-Miatello, *Hypercomplex structures on a class of solvable Lie groups*, Quart. J. Math. Oxford Ser. (2) **47** (1996), no. 1888, 389–404.
- [B1] M. Berger, *Classification des espaces homogènes symétriques irréductibles*, C.R. Acad. Sci. Paris **240** (1955), 2370–2372.
- [B2] M. Berger, *Structure et classification des espaces homogènes symétriques groupe d’isométries semi-simple*, C. R. Acad. Sci. Paris **241** (1955), 1696–1698.
- [C-P] M. Cahen, M. Parker, *Sur des classes d’espaces pseudo-riemanniens symétriques*, Bull. Soc. Math. Bely. **22** (1970), 339–354.
- [C-W] M. Cahen, N. Wallach, *Lorentzian symmetric spaces*, Bull. Amer. Math. Soc. **76** (1970), 585–591.
- [F] A.S. Fedenko, *Spaces defined by Lie group endomorphisms (Φ -spaces)*, Collection of articles dedicated to the memory of German Fedorovič Laptev, Trudy Geometr. Sem. **4** (1973), 231–267.
- [J] D. Joyce, *Compact hypercomplex and quaternionic manifolds*, J. Diff. Geom. **35** (1992), 743–761.
- [O-V] A.L. Onishchik, E.B. Vinberg (Eds.), *Lie groups and Lie algebras III*, Encyclopaedia of Mathematical Sciences **41**, Springer, 1994.

- [P] A.Z. Petrov, *Einstein spaces*, Pergamon, Oxford, 1969.
- [SSTVP] Ph. Spindel, A. Sevrin, W. Troost, A. Van Proeyen *Extended super-symmetric σ -models on group manifolds*, Nuclear Phys. B **308** (1988), 662–698.
- [W] H. Wu, *On the de Rham decomposition theorem*, Illinois J. Math. **8** (1964), 291–311.